Comment on "Unraveling a Clinical Paradox: Why Does Bronchial Thermoplasty Work in Asthma?"

Authors: Bindi S. Brook¹, Igor L. Chernyavsky², Richard J. Russell³, Ruth M. Saunders³ & Christopher E. Brightling³

Affiliations:

¹Nottingham University, Nottingham, NG7 2RD, UK

²University of Manchester, Manchester, M13 9PL, UK

³University of Leicester, LE3 9QP, UK

Originally Published in:

Brook BS, Chernyavsky IL, Russell RJ, Saunders RM & Brightling CE (2019) Comment on "Unraveling a Clinical Paradox: Why Does Bronchial Thermoplasty Work in Asthma?" Am J Respir Cell Mol Biol 61:660–661 (doi:10.1165/rcmb.2019-0095LE).

Copyright © 2019 by the American Thoracic Society. The final publication is available at https://doi.org/10.1165/rcmb.2019-0095LE.

Comment on "Unraveling a Clinical Paradox: Why Does Bronchial Thermoplasty Work in Asthma?"

To the Editor:

We are writing to call for caution when interpreting the predictions of computational models at the organ-level^{1,2} in the context of tissue-scale effects³. Having read recent work by Donovan and colleagues^{1,2} with great interest, we would like to raise a few points about the use of modelling to understand the mechanisms of therapeutic action of bronchial thermoplasty (BT) in asthma control.

First, the authors postulate a fixed 75% airway smooth muscle (ASM) reduction post-BT, quoting published biopsy data^{1,2}. However, relative reductions in ASM mass in reported *in vivo* biopsy studies range from ~ 50-80% as cited and further demonstrated in our study³, and more recently reported by d'Hooghe *et al*⁴. These reductions are also characterised by strong inter- and intrapatient variability, e.g. in our study the observed relative reduction in ASM mass had an interquartile range of 6 to 90%³. Furthermore, acutely, we observed less than 60% reduction in viable ASM cell counts *in vitro* and less than 10% of a large conducting airway was predicted to be heated to therapeutic temperatures *in silico*³. Other works also highlight relative resilience of fibroblasts to thermal injury⁵, in agreement with more ASM cells (structurally similar to fibroblasts) remaining viable at higher temperatures than bronchial epithelial cells³. Even granting possible long-term decrease in ASM content of the treated airways, we still expect a significant degree of heterogeneity. It therefore appears important to identify the threshold percentage in ASM reduction and/or de-activation at the level of individual bronchi that will make a functional impact at the organ level. The future approaches thus lie in integrated tissue- and organ-level models to provide patient-specific spatial heterogeneity of local BT thermal impact.

It is also important to differentiate relative contribution of smaller vs. larger bronchi to the clinical impact of BT (either direct or indirect). The organ-level model¹ suggests the largest airways as a key pathway to improved ventilation. On the other hand, the reactivity of smaller airways is known to be more prominent in heavily remodelled fatal asthma (e.g. [15] in Donovan, *et al.*²), which is confirmed by the results of the study (see Fig. 3(f) in Donovan, *et al.*¹ in agreement with tissue biomechanical models⁶). An integrated tissue model³ also indicates that the potential impact of BT is strongest at the distal end of accessible bronchi and, thus, could affect highly reactive ASM-rich airways in severe near-fatal asthma.

Finally, although the clinical effect of BT on reducing the frequency of exacerbation episodes is better established, the impact of BT mechanisms on lung function appear less clear^{4,7}, with no confirmed correlation between clinical outcomes and improvement in functional respiratory tests. However, there is evidence of BT-induced improvement in epithelial integrity³ that might be associated with long-term altered gene expression profiles in bronchial epithelium⁸, potentially contributing to a lower exacerbation incidence rate.

Bindi S. Brook¹ Igor L. Chernyavsky² Richard J. Russell³ Ruth M. Saunders³ Christopher E. Brightling³ ¹Nottingham University, Nottingham, NG7 2RD, UK ²University of Manchester, Manchester, M13 9PL, UK

References

- Donovan GM, Elliot JG, Green FHY, James AL & Noble PB. Unraveling a Clinical Paradox: Why Does Bronchial Thermoplasty Work in Asthma? *Am J Respir Cell Mol Biol* 2018;59:355-362.
- Donovan GM, Elliot JG, Boser SR, Green FHY, James AL & Noble PB. Patient-specific targeted bronchial thermoplasty: predictions of improved outcomes with structure-guided treatment. J Appl Physiol 2019 Jan. Available from https://doi.org/10.1152/japplphysiol.00951.2018
- 3. Chernyavsky IL, Russell RJ, Saunders RM, Morris GE, Berair R, Singapuri A, Chachi L, Mansur AH, Howarth PH, Dennison P, Chaudhuri R, Bicknell S, Rose FRAJ, Siddiqui S, Brook BS & Brightling CE. In vitro, in silico and in vivo study challenges the impact of bronchial thermoplasty on acute airway smooth muscle mass loss. *Eur Respir J* 2018;51(5) pii: 1701680.
- d'Hooghe JNS, Goorsenberg AWM, Ten Hacken NHT, Weersink EJM, Roelofs JJTH, Mauad T, Shah PL, Annema JT, Bonta PI, TASMA research group. Airway smooth muscle reduction after bronchial thermoplasty in severe asthma correlates with FEV1. *Clin Exp Allergy* 2019 Feb. Available from <u>https://doi.org/10.1111/cea.13365</u>
- 5. O'Neill DP, Peng T, Stiegler P, Mayrhauser U, Koestenbauer S, Tscheliessnigg K & Payne SJ. A three-state mathematical model of hyperthermic cell death. *Ann Biomed Eng* 2011;39:570-579.
- 6. Hiorns JE, Jensen OE & Brook BS. Nonlinear compliance modulates dynamic bronchoconstriction in a multiscale airway model. *Biophysical J* 2014;107:3030-3042.
- Langton D, Ing A, Sha J, Bennetts K, Hersch N, Kwok M, Plummer V, Thien F & Farah C. Measuring the effects of bronchial thermoplasty using oscillometry. *Respirology* 2018 Nov. Available from <u>https://doi.org/10.1111/resp.13439</u>
- 8. Liao SY, Linderholm AL, Yoneda KY, Kenyon NJ & Harper RW. Airway transcriptomic profiling after bronchial thermoplasty. *ERJ Open Res* 2019;5(1) pii: 00123-2018.